

Cosmological evolution of general scalar fields in a brane-world cosmology

Shuntaro Mizuno¹, Seung-Joo Lee², and Edmund. J. Copeland²

¹*Department of Physics, Waseda University, Okubo 3-4-1, Shinjuku, Tokyo 169-8555, Japan and*

²*Department of Physics and Astronomy, University of Sussex, Falmer, Brighton BN1 9QJ, United Kingdom*
(Dated: February 2, 2008)

We study the cosmology of a general scalar field and barotropic fluid during the early stage of a brane-world where the Friedmann constraint is dominated by the square of the energy density. Assuming both the scalar field and fluid are confined to the brane, we find a range of behaviour depending on the form of the potential. Generalising an approach developed for a standard Friedmann cosmology, in [1], we show that the potential dependence $V(\phi)$ can be described through a parameter $\lambda \equiv -\sqrt{2}m_5^{3/2}V'/(\sqrt{H}V)$, where m_5 is the 5-dimensional Planck mass, H is the Hubble parameter and $V' \equiv \frac{dV}{d\phi}$. For the case where λ asymptotes to zero, we show that the solution exhibits stable inflationary behaviour. On the other hand if it approaches a finite constant, then $V(\phi) \propto \frac{1}{\phi^2}$. For $\lambda \rightarrow \infty$ asymptotically, we find examples where it does so both with and without oscillating. In the latter case, the barotropic fluid dominates the scalar field asymptotically. Finally we point out an interesting duality which leads to identical evolution equations in the high energy ρ^2 dominated regime and the low energy ρ dominated regime.

PACS numbers: pacs: 98.80.Cq

I. INTRODUCTION

Scalar fields play very important roles in cosmology, especially in the early stage of the universe. The Inflationary scenario [2] relies on the potential energy of a scalar ‘inflaton’ field to drive a period of early universe acceleration. It is thought that the early universe could well have also been characterised by a series of phase transitions, in which topological defects could have been formed [3, 4]. Moreover in context of the string theory, the natural values of the gauge and gravitational couplings in our 4d universe are explained by the dynamics of ‘moduli’ scalar fields. [5, 6]. More recently, potentials of light scalar fields have been invoked as a possible source of the Dark Energy which appears to be dominating the dynamics of our Universe today. The success of this approach has often been established by demonstrating the existence of attractor solutions by investigating the asymptotic dynamics of the system being studied. For example in the context of usual 4D Einstein gravity, particular potentials have been discussed [7, 8, 9, 10, 11, 12, 13], whereas others have adopted an approach which does not specify a particular potential [1, 14, 15]. However, recent higher-dimensional unification scenarios based on brane-worlds suggest that the gravitational law could be different from Einstein’s during the early stages of our Universe. In the brane-world scenario, our Universe is a four dimensional hypersurface embedded in higher dimensions. Standard-model particles are confined to the brane, while gravity can propagate in the higher-dimensional bulk space [16, 17, 18, 19, 20, 21, 22].

Both the models based on Horava-Witten and Randall-Sundrum II (R-S II) are interesting because they lead to a new type of compactification with gravity. They also make concrete predictions of how conventional gravity will be modified at high-energy scales. Many authors have discussed the cosmology associated with these

scenarios [23, 24, 25, 26, 27, 28, 29]. When we consider the homogeneous and isotropic cosmology based on these brane world models, the difference from conventional cosmology can be related to the appearance of two new terms in the Friedmann equation, i.e., the quadratic term of the energy-momentum and a dark radiation term [30, 31, 32]. The allowed impact of dark radiation, is constrained by nucleosynthesis [33], with the most important change in the scalar field dynamics being due to the appearance of the quadratic energy density term.

In this paper, we study the dynamics of a scalar field which is confined to the brane in the context of a R-S II brane-world. As well as the scalar field, we include a barotropic fluid on the brane (which could be radiation or matter). Since the modification of the cosmic expansion law arising through the quadratic energy density term becomes important during the early stage of the universe, we concentrate on this stage. In order to classify the asymptotic behaviour of the solutions, we follow the model independent approach as first proposed in [1, 14]. Earlier work investigating the impact of these ρ^2 corrections has concentrated on models with particular potentials [34, 35, 36, 37, 38, 39, 40]. We intend to go beyond that in this paper, and in passing note that the analysis complements our previous discussion on scaling solutions in ρ^2 cosmologies [41].

The organization of the paper is as follows. In Sec. II, we review the key equations of motion and define variables which will allow us to analyse the asymptotic behaviour of this system and point out an intriguing duality with the equations obtained in the context of scaling solutions in standard Friedman cosmologies [1, 11]. In Sec. III we obtain the class of attractor solutions which exist for constant λ . We then extend the analysis to the cases where λ evolves and find classes of solutions corresponding to inflationary regimes (Sec. IV); scalar field kinetic energy dominated regimes (Sec. V), and oscillating

regimes (Sec. VI). We summarise our results in Section VII.

II. EQUATIONS OF MOTION

We analyze the dynamics of a scalar field in a Randall-Sundrum II (R-S II) brane-world [22], because the model is simple and concrete. However, we expect our key results to also hold in other brane-world models, in which a quadratic term in the energy-momentum tensor generically appears. In the R-S II model, even though the extra-dimension is not compactified, gravity is confined to the brane at low energy, resulting in Newtonian gravity in our world which is described by the intrinsic metric of the 4-dimensional brane spacetime. By use of Israel's junction condition and assuming Z_2 symmetry, the gravitational equations on the 3-brane are given by [30]

$$\begin{aligned} {}^{(4)}G_{\mu\nu} &= -{}^{(4)}\Lambda g_{\mu\nu} + \kappa_4^2 T_{\mu\nu} + \kappa_5^4 \Pi_{\mu\nu} - E_{\mu\nu}, \\ \Pi_{\mu\nu} &\equiv -\frac{1}{4}T_{\mu\alpha}T_{\nu}^{\alpha} + \frac{1}{12}TT_{\mu\nu} + \frac{1}{8}g_{\mu\nu}T_{\alpha\beta}T^{\alpha\beta} - \frac{1}{24}g_{\mu\nu}T^2, \end{aligned} \quad (1)$$

where ${}^{(4)}G_{\mu\nu}$ is the Einstein tensor with respect to the intrinsic metric $g_{\mu\nu}$, ${}^{(4)}\Lambda$ is the 4-dimensional cosmological constant, $T_{\mu\nu}$ represents the energy-momentum tensor of matter fields confined to the brane and $\Pi_{\mu\nu}$ is the local correction term as a result of a brane embedded in the bulk. $E_{\mu\nu}$ is a part of the 5-dimensional Weyl tensor and carries information about the bulk geometry. $\kappa_4^2 = 8\pi G_4$ and $\kappa_5^2 = 8\pi G_5$ are 4-dimensional and 5-dimensional gravitational constants, respectively. In what follows, we use the 4-dimensional Planck mass $m_4 \equiv \kappa_4^{-1} = (2.4 \times 10^{18} \text{ GeV})$ and the 5-dimensional Planck mass $m_5 \equiv \kappa_5^{-2/3}$. m_5 is related to m_4 in terms of the brane tension σ as $\sigma m_4^2 = 6m_5^6$, and could be much smaller than m_4 .

Assuming the Friedmann-Robertson-Walker spacetime in our brane world, we find the effective Friedmann equations arising from Eq.(1) are

$$\begin{aligned} H^2 + \frac{k}{a^2} &= \frac{1}{3}{}^{(4)}\Lambda + \frac{1}{3m_4^2}\rho + \frac{1}{36m_5^6}\rho^2 + \frac{\mathcal{C}}{a^4} \\ \dot{H} - \frac{k}{a^2} &= -\frac{1}{2m_4^2}(P + \rho) - \frac{1}{12m_5^6}\rho(P + \rho) - \frac{2\mathcal{C}}{a^4} \end{aligned} \quad (2)$$

where a is a scale factor of the universe, $H = \dot{a}/a$ is its Hubble parameter, k is a curvature constant, P and ρ are the total pressure and energy density of matter fields, respectively. $\mathcal{C}$ is a constant related to “dark radiation” coming from $E_{\mu\nu}$ [31, 32]. In what follows, we consider only a flat Friedmann model ($k = 0$). We also assume that ${}^{(4)}\Lambda$ vanishes at least in the early stage of the universe and $\mathcal{C}$ is negligible since it should be diluted by inflation and is strongly constrained at the time of Nucleosynthesis [33]. (However, it must be noted that if we consider perturbations around the back ground solutions

in this paper, we must consider the term arising from $E_{\mu\nu}$.)

For matter fields on the brane, we consider a scalar field ϕ with a potential $V(\phi)$ as well as a barotropic fluid with an equation of state $P_B = (\gamma - 1)\rho_B$, where γ is an adiabatic index. (We will mainly be considering the case of radiation with $\gamma = 4/3$, since we are interested in the early stage of the universe before nucleosynthesis.) We will also assume that the two fluids do not couple to each other explicitly.

Although a 5-dimensional scalar field living in the bulk [43] may also appear in other brane-world scenarios, in this paper we only consider a 4-dimensional scalar field confined to the brane. Such a scalar field might originate in the condensation of fermionic matter fields confined to the brane.

As the universe expands, the energy density decreases. This means that the quadratic term could have been very important in the early stages of the universe. Comparing the energy density terms in Eq. (2), we find that the quadratic term dominates when

$$\rho > \rho_c \equiv 12m_5^6/m_4^2. \quad (4)$$

In this regime the expansion law of the Universe is clearly modified. For example, the expansion law in the radiation-dominant era becomes $a \propto t^{1/4}$ compared to the conventional case, $a \propto t^{1/2}$. If such a period of ρ^2 domination did exist, then Nucleosynthesis would have provided us with a constraint on m_5 , since it must have occurred during the conventional-radiation dominated era in order to explain the abundances of the light elements. For example, assuming that the energy density at $a = a_c$ is dominated by radiation as $\rho_c \sim (\pi^2/30)gT_c^4$, where g is the number of degrees of freedom of light particles, the temperature of the universe T_c must be higher than that of nucleosynthesis, i.e. $T_c > T_{NS} \sim 1 \text{ MeV}$. This constraint implies that $m_5 > 1.6 \times 10^4 (g/100)^{1/6} (T_{NS}/1 \text{ MeV})^{2/3} \text{ GeV}$. (For the brane tension σ , this constraint is $\sigma^{1/4} > 2.0 \text{ MeV}$).

In the rest of the paper we will be concentrating on the evolution of the universe in the regime dominated by the ρ^2 contribution to the energy density. We will find there is a way of performing this analysis in an analogous manner to that already developed for the usual cosmologies dominated by the linear ρ term [1, 11, 14]. In fact we will introduce a set of variables which will lead to exactly the same form of evolution equations for the two regimes, allowing us to relate the attractor solutions that exist in the two regimes through duality transformations. Of course, in a realistic cosmology, the ρ^2 dominated regime is eventually replaced by the linear regime. Any asymptotic solution in the ρ^2 regime should be reached before the linear regime takes over at $\rho \sim \rho_c$, and we confirm that is the case in this analysis.

In the regime where $\rho \gg \rho_c$, Eq. (2) along with conservation of energy and momentum for each fluid component

on the brane gives

$$\dot{H} = -\frac{1}{12m_5^6}(\rho_B + \frac{1}{2}\dot{\phi}^2 + V)(\gamma\rho_B + \dot{\phi}^2), \quad (5)$$

$$\dot{\rho}_B = -3\gamma H\rho_B, \quad (6)$$

$$\ddot{\phi} = -3H\dot{\phi} - \frac{dV}{d\phi}, \quad (7)$$

which are subject to the Friedmann constraint given in Eq. (2)

$$H = \frac{1}{6m_5^3}(\rho_B + \frac{1}{2}\dot{\phi}^2 + V), \quad (8)$$

where dots denote derivatives with respect to time. The energy density and pressure of the homogeneous scalar field are given by $\rho_\phi = \dot{\phi}^2/2 + V$ and $P_\phi = \dot{\phi}^2/2 - V$, respectively. The effective adiabatic index of the scalar field at any time,

$$\gamma_\phi \equiv \frac{\rho_\phi + P_\phi}{\rho_\phi} = \frac{\dot{\phi}^2}{\dot{\phi}^2/2 + V}, \quad (9)$$

is, in general, time or scale dependent.

In order to analyze the stability of the solutions, we modify the variables, $X_{\text{CLW}}, Y_{\text{CLW}}$ first introduced in Ref. [11] to investigate the evolution of a scalar field in a conventional Friedmann cosmology. Defining

$$X = \frac{m_5^{-3/2}}{2\sqrt{3}} \frac{\dot{\phi}}{\sqrt{H}}, \quad (10)$$

$$Y = \frac{m_5^{-3/2}}{\sqrt{6}} \frac{\sqrt{V}}{\sqrt{H}}, \quad (11)$$

where a prime denotes a derivative with respect to the logarithm of a scale factor a , $N \equiv \ln a$, we see that these are related to those defined in Ref. [11] by

$$\frac{X}{X_{\text{CLW}}} = \frac{Y}{Y_{\text{CLW}}} = \frac{\kappa_5}{\kappa_4} \sqrt{\frac{H}{2}} \quad (12)$$

$$(13)$$

In terms of these new variables, the equations of motion read,

$$X' = -3X + \lambda\sqrt{\frac{3}{2}}Y^2 + \frac{3}{2}X[2X^2 + \gamma(1 - X^2 - Y^2)], \quad (14)$$

$$Y' = -\lambda\sqrt{\frac{3}{2}}XY + \frac{3}{2}Y[2X^2 + \gamma(1 - X^2 - Y^2)], \quad (15)$$

$$\lambda' = -\sqrt{6}\lambda^2(\Gamma - 1)X + \frac{3}{2}\lambda[2X^2 + \gamma(1 - X^2 - Y^2)], \quad (16)$$

where generalising the expressions in Refs. [1] and [8], we have defined

$$\lambda \equiv -\sqrt{2}\frac{m_5^{3/2}}{\sqrt{H}}\frac{dV/d\phi}{V}, \quad (17)$$

$$\Gamma \equiv V \frac{d^2V/d\phi^2}{(dV/d\phi)^2}. \quad (18)$$

Following Ref. [1], we see that the equation for H can also be written down as

$$H' = -3H[2X^2 + \gamma(1 - X^2 - Y^2)]. \quad (19)$$

Since the energy density of the barotropic fluid satisfies ($\rho_B \geq 0$), the contribution of the scalar field to the total energy density, $\Omega_\phi \equiv \rho_\phi/(6m_5^3H) = X^2 + Y^2$ is bounded, $0 \leq X^2 + Y^2 \leq 1$. Hence the evolution of this system is completely described by trajectories within the unit disk. Moreover, since the system is symmetric under the reflection $(X, Y) \rightarrow (X, -Y)$ and the evolution will not go beyond the $Y = 0$ line (which corresponds to $\phi = \infty$ or $H = \infty$), it is enough to discuss the upper half disk ($Y \geq 0$).

There is an intriguing aspect to these evolution equations. Given the definitions of X and Y above, Eqs. (14-16) are identical in form to those for $X_{\text{CLW}}, Y_{\text{CLW}}$ and $\lambda_{\text{CLW}} \equiv -\frac{1}{\kappa_4}\frac{V'}{V}$ where λ and λ_{CLW} are related through

$$\frac{\lambda}{\lambda_{\text{CLW}}} = \frac{\kappa_4}{\kappa_5} \sqrt{\frac{2}{H}} \quad (20)$$

in [1, 11, 14]. This means that the attractor solutions for X_{CLW} etc, obtained in the standard Friedmann case where the Hubble parameter is determined by the linear energy density term, are the same attractor solutions in terms of our new variables X etc... but for the case of ρ^2 domination. Even though they do not correspond to the same potential or evolution of the Hubble parameter, they do correspond to the same parameters in terms of X, Y and λ . This is intriguing, as it implies there is a duality between the high energy ρ^2 set of solutions and the low energy solutions corresponding to ρ domination.

III. ATTRACTOR SOLUTIONS

We want to now turn our attention to the evolution equations (14) - (16). We are particularly interested in the case of scaling solutions, where $X' = Y' = \lambda' = 0$. To start with we consider the case where λ tends to a constant value asymptotically. Using Eq. (15), for non-trivial X, Y and λ , we see that the condition $\lambda' = 0$ in Eq. (16) corresponds to the constraint $\Gamma = \frac{3}{2}$. Integrating, Eq. (18) this in turn implies the form for the potential which leads to a constant λ , namely

$$V(\phi) = \frac{\mu^6}{\phi^2} \quad (21)$$

where μ is a constant with the dimensions of mass. This result was first obtained using a different method in [41]. The equivalent potential which leads to scaling

solutions with constant λ_{CLW} in the case of the standard Friedmann cosmology is an exponential potential $V(\phi) = V_0 \exp(-\lambda_{\text{CLW}} \kappa_4 \phi)$ [11], which is clear from Eqs. (17) and (20). We can now summarise the attractor solutions X_c, Y_c , for our scaling potential Eq. (21), recalling that they correspond to the same scaling solutions $X_{\text{CLW}}, Y_{\text{CLW}}$ as found in [11]. Eqs. (14) and (15) admit five different critical solutions for X, Y with λ constant.

For these critical solutions, $(X_c = 1, Y_c = 0), (X_c = -1, Y_c = 0), (X_c = 0, Y_c = 0)$ are unstable (extreme) critical points. The other two solutions depend on the value of λ . For $\lambda^2 > 3\gamma$, we find the critical attractor values

$$\begin{aligned} X_c &= \sqrt{\frac{3\gamma}{2\lambda}}, \\ Y_c &= \sqrt{\frac{3(2-\gamma)\gamma}{2\lambda^2}}, \end{aligned} \quad (22)$$

which satisfies $\gamma_\phi \equiv \frac{2X^2}{X^2+Y^2} = \gamma$ implying that both the barotropic fluid and scalar field energy densities scale with the same redshift. The fractional energy density stored in the scalar field is given by $\Omega_{\phi c} = (3\gamma)/\lambda^2$. If we consider the case of radiation for the barotropic fluid, then from Eqs. (11), (17) and (22) we obtain $\Omega_{\phi c} = (1/2)(\mu/m_5)^3$ [41]. It is worth noting that because of the nature of the attractor solutions, for $3\gamma < \lambda^2 < 6\gamma$, even if the scalar field dominates the universe, the cosmic expansion law becomes the same as the barotropic fluid dominated universe.

If the asymptotic behavior of this system is characterized by this scaling solution, then once the energy density has decreased to $\rho \sim \rho_c$, we recover Einstein gravity with the linear energy density term dominating over the quadratic term in Eqs. (2) and (3). It is well known that in this conventional cosmology, a scalar field with potential $V = \mu^6 \phi^{-2}$ leads to a tracker solution and dominates the universe at late times [7, 8]. In order for this term to then provide the observed Dark Energy today, it requires μ to be fine tuned to a value $\text{O}(\text{GeV})$, meaning that $\Omega_{\phi c}$ during the period that the quadratic term dominates must be extremely small.

On the other hand, if $\lambda^2 < 6$, the corresponding attractor solution is

$$\begin{aligned} X_c &= \frac{\lambda}{\sqrt{6}}, \\ Y_c &= \sqrt{1 - \frac{\lambda^2}{6}}, \end{aligned} \quad (23)$$

which satisfies $\gamma_\phi = (1/3)\lambda^2$ and $\Omega_{\phi c} = 1$. In this case the scalar field energy density dominates the universe asymptotically. If the scalar field has $\gamma_\phi < \gamma$, then the solutions in Eqs. (23) are stable, and the redshift of the scalar field is slower than that of the barotropic fluid. It corresponds to power-law inflation and is stable as in the conventional cosmology driven by an exponential potential [11, 45]. Even though this solution is interesting

because it is exact, it has the drawback of failing to stop inflating, so in a successful cosmology it would have to eventually end inflation and recover the conventional cosmology. Furthermore, for such a large value of μ , this scalar field dominates the universe much earlier. However, if $\gamma_\phi = (1/3)\lambda^2 > \gamma$, then the solutions in Eqs. (23) are unstable and the scalar field ends up in the regime of the solution given in Eqs. (22).

Having considered the case of a constant λ , we now turn our attention to the case where $\lambda(N)$ is not constant. Then the critical values in Eqs. (22) and (23) hold $X' = Y' = 0$ only at single points, not over intervals of time. This means that the attractor solutions to Eqs. (14) and (15) are only valid as an asymptotic limit and in general X_c, Y_c, λ are time dependent. If X, Y do not oscillate, then since their values are constrained to $|X| \leq 1, |Y| \leq 1$, this implies that they will approach constant values asymptotically, given by the attractor solutions of Eqs. (14) and (15), and X', Y' will vanish. Therefore, we can generalize the attractor solutions of X, Y given in Eqs. (22) and (23) for more complicated potentials that have a nonconstant $\lambda(N)$. In the following sections, we will analyze other asymptotic solutions making use of this generalization.

IV. POTENTIAL DOMINATED INFLATIONARY SOLUTIONS

We now consider the case where $\lambda = -\frac{\sqrt{2}m_5^{3/2}}{\sqrt{H}} \frac{V'}{V} \rightarrow 0$, asymptotically. We might naively expect this happens as long as $V' \rightarrow 0$ faster than V as $\phi \rightarrow \infty$, such as the case for inverse power law potentials of the form $V(\phi) = \mu^{4+n}\phi^{-n}$, with $n > 0$. Indeed, in the conventional Friedmann cosmology in which $\lambda_{\text{CLW}} \equiv -\frac{1}{\kappa_4} \frac{V'}{V}$ plays the same role as our λ , our expectation is realised in that $\lambda_{\text{CLW}} \rightarrow 0$, asymptotically [1]. However in our case, if we substitute $V(\phi) = \mu^{4+n}\phi^{-n}$ with $n > 0$ into Eq. (17), and use the fact that in the regime where the potential is dominating, $H \propto V$, we obtain $\lambda \propto \phi^{\frac{n}{2}-1}$. From this, even with inverse power law potentials, only when $2 > n > 0$, does $\lambda \rightarrow 0$ as $\phi \rightarrow \infty$.

In the following we will concentrate on models with potentials which lead to $\lambda \rightarrow 0$ asymptotically. If this is realized, we can eliminate the term proportional to λ in Eq. (14), and Eq. (15). After rewriting them in terms of H' with Eq. (19), and since $-6 < H'/H < 0$ for all values of X, Y , and γ , we obtain the following relations identical to those obtained in [1], but recalling the interpretation for the variables X, Y and λ differ in the two cases:

$$\begin{aligned} \frac{X'}{X} &= -\left(3 + \frac{1}{2} \frac{H'}{H}\right) < 0, \\ \frac{Y'}{Y} &= -\frac{1}{2} \frac{H'}{H} > 0. \end{aligned} \quad (24)$$

This of course is to be expected given the duality we discussed earlier. As explained in [1] these equations in-

indicate that $X \rightarrow 0$ its minimum value, as $Y \rightarrow 1$, its maximum value. This allows us to solve Eq. (14), Eq. (15), and Eq. (19) for X , Y , and H in the region $|X| \ll 1$, $|\lambda| \ll 1$:

$$\begin{aligned} X(N) &= \frac{e^{-3N}}{\sqrt{1 - ce^{-3\gamma N}}}, \\ Y(N) &= \frac{1}{\sqrt{1 - ce^{-3\gamma N}}}, \\ H(N) &= d(1 - e^{-3\gamma N}), \end{aligned} \quad (25)$$

where c and d are integration constants. These approximate solutions are consistent with those in Eq. (23) in the limit $\lambda \rightarrow 0$, whilst recalling that X and Y are varying with time. They show that the scalar field potential dominates the energy density of the universe as the universe inflates with almost constant Hubble parameter. In fig. 1, we demonstrate the validity of the solutions in Eq. (25), as we show numerical solutions for a model with a barotropic fluid of radiation ($\gamma = 4/3$), and with potential $V(\phi) = \mu^5 \phi^{-1}$.




FIG. 1: Evolution of a Universe driven by the ρ^2 term in the Friedmann equation. The universe contains radiation and a scalar field with $V = \mu^5 \phi^{-1}$. For simplicity, we choose $\mu = m_5$. The initial conditions are $X_0 = 0.2$, $Y_0 = 0.4$. In (a) the Hubble parameter (solid curve) tends to a finite constant leading to inflation. In (b) we show Ω_ϕ quickly approaching unity. In (c) λ evolves to zero asymptotically. In (d), the numerical solutions for Y (upper curve) and X (lower curve) are plotted. The solutions given by Eqs. (25) are obtained asymptotically.

This asymptotically inflating solution is stable, hence in order to recover standard cosmology at late times, a mechanism to end this period of inflation is necessary. Simply relying on the existence of the ρ term dominating in the Friedmann equation will not be sufficient here,

because as shown in [1], this potential also has an inflationary solution in that era. This can be readily seen from Eq. (20), where $\lambda_{\text{CLW}} = (\kappa_5/\kappa_4)\sqrt{H/2}\lambda \rightarrow 0$ as $\lambda \rightarrow 0$ for constant H .

V. KINETIC-TERM DOMINATED SOLUTION

Having considered the case $\lambda \rightarrow \text{const}$ (including zero), we now consider the case where asymptotically $\lambda = -\frac{\sqrt{2}m_5^{3/2}}{\sqrt{H}}\frac{V'}{V} \rightarrow \infty$ smoothly without oscillations.

The presence of the $1/\sqrt{H}$ term in the definition of λ , as opposed to the definition of $\lambda_{\text{CLW}} \propto V'/V$, and the fact that in most cosmologies $H \rightarrow 0$ asymptotically, means that a wider class of potentials satisfy $\lambda \rightarrow \infty$, than satisfy $\lambda_{\text{CLW}} \rightarrow \infty$ [1]. For example, as we show below, the inverse polynomial potentials, $V(\phi) = \mu^{4+n}\phi^{-n}$ with $n > 2$ discussed earlier satisfy this condition. Another concrete example is a model including an exponential potential which satisfies $\lambda \propto 1/\sqrt{H} \rightarrow \infty$.

Although the specific potentials included in this limit may be different between the ρ^2 and ρ cosmologies, the invariance of the form of the equations of motion (14)-(15) when written in terms of X and Y implies that the solutions obtained in terms of X and Y in Ref. [1] for the case of the conventional ρ dominated cosmology, apply to our case as well. We therefore summarise the results presented there.

Initially we expect X and Y to be order unity, which means for $|\lambda| \gg 1$, the leading terms of Eq. (14) and Eq. (15) are

$$\begin{aligned} X' &= \sqrt{\frac{3}{2}}\lambda Y^2, \\ Y' &= -\sqrt{\frac{3}{2}}\lambda XY. \end{aligned} \quad (26)$$

Note for positive (negative) λ , $X \rightarrow 1(-1)$ whereas for both signs, $Y \rightarrow 0$. Eventually as Y keeps decreasing, other terms in Eqs. (14) and (15) become important. In particular if $|X'| \geq |\lambda|Y^2$, then using Eq. (19), the evolution of X is given by

$$\frac{X'}{X} = -\left(3 + \frac{H'}{2H}\right). \quad (27)$$

Now since $-6 < H'/H < 0$, X having reached a maximum value turns over and like Y , heads off finally approaching the values given by the solution of Eqs. (22), $X \rightarrow X_c$, $Y \rightarrow Y_c$. For X_c and Y_c with $|\lambda| \gg 1$, then $1 \gg |X_c| > Y_c$ holds, and ρ_ϕ will decrease faster than ρ_B with $\Omega_\phi \rightarrow 0$. Even though X_c , Y_c are not critical (constant) points since λ is not constant, we have verified numerically that the above asymptotic solutions are good approximations. In Fig. 2, typical examples of the numerical results for a model with potential $V(\phi) = \mu^7 \phi^{-3}$ are shown, in a universe containing radiation. Note in particular the behaviour of $\lambda \rightarrow \infty$ in (c) and how in (d)

X increases initially before decreasing towards zero, approaching the solutions given in Eqs. (22). As expected these solutions match closely those presented in Figure 2 of Ref [1], for the conventional ρ dominated cosmology, but there are important differences. In particular the form of the potential involved (exponential in their case, inverse polynomial in ours), and most significantly the fact we are dealing with different cosmologies.




FIG. 2: Evolution of the ρ^2 cosmology for a universe filled with radiation and a scalar field with $V = \mu^7 \phi^{-3}$. For simplicity, we choose $\mu = m_5$. The initial conditions are $X_0 = 0.2$, $Y_0 = 0.6$. In (a) we show that the Hubble parameter normalized by its initial value (solid curve) tends to zero. In (b) we show Ω_ϕ quickly approaching 0. In (c) λ evolves to infinity asymptotically. In (d), the numerical X (solid curve) and Y (dotted curve) solutions are plotted. As in Eqs. (26), at the beginning, X increases and reaches its maximum value, before turning over following Eq. (27) as it (and Y) approaches 0 asymptotically.

Let us briefly digress to discuss the impact of this sort of solution. We have seen that as long as $\mu \leq m_5$, then the asymptotic behaviour in the ρ^2 cosmology, is one where $\rho_\gamma > \rho_\phi$. This will carry on until $\rho \sim \rho_c$, after which we will enter a conventional radiation dominated universe. This is a nice feature as once in that regime we know from earlier studies that the inverse polynomial potentials being discussed here lead to tracking behaviour necessary to explain the dark energy today. The power of the ρ^2 regime here lies in the fact that it provides a dynamical explanation of the large difference between the energy density of the dominant barotropic fluid and the scalar field ‘initially’ as we enter the conventional cosmology [42].

If we decide to concentrate initially on the limit $\phi \rightarrow 0$, then this corresponds to $\lambda \rightarrow 0$ suggesting an unstable

inflationary solution in this limit, for the inverse polynomial potentials $V(\phi) = \mu^{4+n} \phi^{-n}$, with $n > 2$. This result provides one of a number of possibilities which have been investigated where the same scalar field potential is used to realize both a period of early inflation and late time dark energy domination, although there are strong constraints on their viability arising from the tendency to overproduce gravitational waves during reheating. This is a form of quintessential inflation [46], except that it is making use of the brane-world scenario, and has been investigated recently by a number of authors [34, 35, 36, 37, 38, 39, 40, 41].

VI. OSCILLATING SOLUTIONS

We now turn our attention to consider the case $|\lambda| \rightarrow \infty$ with the scalar field ϕ oscillating about the minimum of its potential asymptotically. In the context of realistic scenarios this is somewhat of a formal exercise in that we might well expect that the $\rho \gg \rho_c$ constraint would have been violated once this situation had been reached. However, we feel that it is worth investigating in its own right as it allows us to pursue the duality relation we have identified in the paper between the two regimes of ρ domination and ρ^2 domination.

Given that ϕ is oscillating about its minimum, we can without loss of generality, take it to be zero, and expand $V(\phi)$ as a power series about the minimum. Keeping only the leading term, we have $(1/n)\mu^{4-n}\phi^n$, where $n = 2, 4, 6, \dots$, even because of the boundedness of the potential. (For $n = 4$ we use $V(\phi) = (1/4)\hat{\lambda}\phi^4$ where $\hat{\lambda}$ is a dimensionless constant.)

Again because of the duality invariance, the equations match those in [1], so we follow their path to determine under which conditions, Ω_ϕ : either dominates (goes to unity); oscillates around a finite constant value; or vanishes asymptotically. In actual fact in [41], we have already analyzed the cases for $n = 2$ and 4 obtaining analytic solutions by making assumptions about which components of the energy density dominate the universe, initially. In these cases, we have obtained approximate solutions by invoking the virial theorem, in which the relation between the time-averaged value of the kinetic term and the potential term of the energy density of the scalar field is given as

$$\langle \dot{\phi}^2 \rangle = n \langle V(\phi) \rangle. \quad (28)$$

From Eq. (28), it is clear that the scalar field behaves as a pressureless perfect fluid (dust fluid) ($\langle \gamma_\phi \rangle = 1$) for $n = 2$, while it behaves as a radiation fluid ($\langle \gamma_\phi \rangle = 4/3$) for $n = 4$. Therefore, if we consider the case of radiation for the barotropic fluid in the early universe, for the case $n = 2$, then the energy density in the scalar field will decrease slower than that of radiation, leading to $\Omega_\phi \rightarrow 1$. For the $n = 4$, case, both fluids evolve at the same rate, hence $\Omega_\phi \rightarrow cte$, a constant value determined by the self coupling parameter $\hat{\lambda}$ in $V(\phi)$.

In order to analyze the case for general n , and without specifying which component dominates the universe, we will introduce the total adiabatic index γ_{tot} . When it is the barotropic fluid that dominates, then $\gamma_{\text{tot}} = \gamma$, but when the scalar field dominates the universe, we take $\gamma_{\text{tot}} = \langle \gamma_\phi \rangle$ since in this case γ_ϕ oscillates. Now, in terms of γ_{tot} , $H(t) = 1/(3\gamma_{\text{tot}}t)$, hence in the ρ^2 dominated era, Eq. (7) can be rewritten as,

$$\ddot{\phi} + \frac{1}{\gamma_{\text{tot}}t} \dot{\phi} + \mu^{4-n} \phi^{n-1} = 0. \quad (29)$$

(Recall for the case $n = 4$, the third term must be $\hat{\lambda}\phi^3$. In that case, if we introduce a new time variable $\tau \equiv \sqrt{\hat{\lambda}}t$, the differential equation takes the same form as Eq. (29) with $n = 4$. Where appropriate we use this time variable for the case $n = 4$.)

Even though we have already obtained the results for the case $n = 2$ in [41], we first consider this case to confirm our previous results. For $V(\phi) = (1/2)\mu^2\phi^2$, the solution of Eq. (29) is given in terms of J_m and Y_m , the Bessel functions of the first and second kind, respectively,

$$\begin{aligned} \phi(z) &= \mu z^m [c_1 J_m(z) + c_2 Y_m(z)], \\ \dot{\phi}(z) &= \mu^2 z^m [c_1 J_{m-1}(z) + c_2 Y_{m-1}(z)], \end{aligned} \quad (30)$$

where c_1 , and c_2 are constants, $m \equiv (1/2) - (1/2\gamma_{\text{tot}})$, $z \equiv \mu t$ [12]. From Eq. (30), X and Y are expressed as

$$\begin{aligned} X &= r z^{m+\frac{1}{2}} [kJ_{m-1}(z) + Y_{m-1}(z)], \\ Y &= r z^{m+\frac{1}{2}} [kJ_m(z) + Y_m(z)], \end{aligned} \quad (31)$$

where $r \equiv (1/2)(\mu/m_5)^{3/2} \sqrt{\gamma_{\text{tot}}} c_2$, $k \equiv c_1/c_2$. A simple analytic expression can be obtained using the formula for the asymptotic limit of the Bessel functions $J_m \sim \sqrt{(2/\pi z)} \cos[z - \pi(2m+1)/4]$, $Y_m \sim \sqrt{(2/\pi z)} \sin[z - \pi(2m+1)/4]$ for $z \gg 1$. The amplitude of X and Y in Eqs. (31) in the limit $z \gg 1$ goes as $X \sim Y \sim z^m$. A finite value of X , Y requires $m = (1/2) - (1/2\gamma_{\text{tot}}) = 0$ (i.e. $\gamma_{\text{tot}} = 1$). Next, we analyze how the asymptotic value of γ_{tot} are decided. In this case, regardless of the form of the barotropic fluid, in the limit $z \gg 1$, $\langle X^2 \rangle = \langle Y^2 \rangle = (r^2/\pi) z^{2m} (k^2 + 1)$, which yields $\langle \gamma_\phi \rangle = 1$. This is completely consistent with the previous results relying on the virial theorem. We can confirm, therefore, if the barotropic fluid has $\gamma < 1$, i.e. smaller than γ_ϕ , then the barotropic fluid decays slower than the scalar field, which yields $\gamma_{\text{tot}} < 1$. In this case, from the asymptotic limit of the Bessel functions, $\Omega_\phi = X^2 + Y^2 \rightarrow 0$. On the other hand, if the barotropic fluid has $\gamma > 1$, i.e. greater than γ_ϕ , then the barotropic fluid decays faster than the scalar field. In this case, $\gamma_{\text{tot}} \rightarrow \langle \gamma_\phi \rangle = 1$, which leads to $\Omega_\phi = X^2 + Y^2 \rightarrow 1$ asymptotically. For the case $\gamma = 1 = \gamma_{\text{tot}}$, Ω_ϕ asymptotes to some constant value between 0 and 1.

We have obtained an analytic solution of Eq. (29) for the case $V(\phi) = (1/2)\mu^2\phi^2$. This is clearly the simplest

case since Eq. (29) is linear in ϕ and its derivatives. For $V(\phi) = (1/n)\mu^{4-n}\phi^n$ with $n > 2$, Eq. (29) becomes non-linear in ϕ and no simple analytic solution exists. However, using the following ansatz,

$$\phi = \mu z^{-\frac{1}{n}} \left[c_1 \cos \left(\beta z^{\frac{(n+2)}{2n}} \right) + c_2 \sin \left(\beta z^{\frac{(n+2)}{2n}} \right) \right], \quad (32)$$

we obtain a solution with the correct asymptotic behavior $\phi^{n/2}$, $\dot{\phi} \propto t^{-1/2}$, $H \propto t^{-1}$, and $\ddot{\phi} \propto \phi^{n-1}$, necessary if we wish X , Y to have finite values other than 0 and 1. The $z(= \mu t)$ exponents in Eq. (32) are such that Eq. (29) is satisfied up to leading orders $t^{-3/2}$ and $t^{(1-n)/n}$. The same condition also imposes the conditions $\gamma_{\text{tot}} = 2n/(2+n)$ and $\beta^2 = (2n/n+2)^2 c_2^{n-2} [\sin(\beta z^{2/n}) + k \cos(\beta z^{2/n})]^{n-2}$. Notice that the value of γ_{tot} is precisely the one we obtained earlier based on general arguments [cf. Eq. (28)]. Of course, the ansatz in Eq. (32) is not a complete solution to Eq. (29) as can be seen in the fact that β is not a true constant (except for the case $n = 2$). For $n > 2$ we must use the average value of β evaluated in the asymptotic region. In terms of Eq. (32), X and Y take the following expressions:

$$\begin{aligned} X &= X_0 \left[\cos \left(\beta z^{\frac{(n+2)}{2n}} \right) - k \sin \left(\beta z^{\frac{(n+2)}{2n}} \right) \right], \\ Y &= Y_0 \left[\sin \left(\beta z^{\frac{(n+2)}{2n}} \right) + k \cos \left(\beta z^{\frac{(n+2)}{2n}} \right) \right]^{\frac{n}{2}}, \end{aligned} \quad (33)$$

with $X_0 = \frac{(n+2)}{4n} \sqrt{\gamma_{\text{tot}}} \beta (\mu/m_5)^{3/2} c_2$ and $Y_0 = \sqrt{\gamma_{\text{tot}}/2n} (\mu/m_5)^{3/2} c_2^{n/2}$.

From Eq. (33) we obtain $\langle Y^2 \rangle / \langle X^2 \rangle = 2/n$, asymptotically, as in Eq. (28), which is consistent with the virial theorem. We have, therefore, $\langle \gamma_\phi \rangle = 2n/(2+n)$ and $\gamma_{\text{tot}} = \langle \gamma_\phi \rangle$, i.e., the ansatz in Eq. (32) is a solution to Eq. (29) only when the dominant energy density redshifts as fast as the scalar field. This is, of course, no surprise since we imposed on the ansatz Eq. (32), the limit $X, Y \rightarrow cte$ ($\neq 0$ or 1).

Even though the above discussion strictly relies on the time-averaged value, we have confirmed that it provides a good approximation to the true numerical results. In fig. 3, typical examples of the numerical results for models with $V(\phi) = (1/2)\mu^2\phi^2$ (in (a) and (b)) and $V(\phi) = (1/4)\lambda\phi^4$ (in (c) and (d)) are plotted. As for the barotropic fluid, we use radiation ($\gamma = 4/3$). For both potentials, we draw the time evolution of Ω_ϕ and Ht .

To conclude this part of the analysis, if the initially dominant energy density component has a larger (smaller) adiabatic index than $\langle \gamma_\phi \rangle = 2n/(2+n)$, then Ω_ϕ will approach 1 (0). For example, if we consider radiation as the barotropic fluid, then for $n > 4$, the energy density of the scalar field will decrease faster than radiation, which guarantees the recovery of the standard cosmology at an epoch when the energy density is lower than ρ_c .

Even though the form of these relations are the same as in a conventional cosmology, the oscillating behavior found in the regime dominated by the quadratic term




FIG. 3: Evolution of the universe filled with radiation and a scalar field with potential $V = (1/2)\mu^2\phi^2$ ((a), (b)) and $V = (1/4)\lambda\phi^4$ ((c), (d)) in the ρ^2 dominating era. For simplicity, we choose $\mu = m_5$ and $\lambda = 1$. The initial conditions are $X_0 = 0.1$, $Y_0 = 0.5$. We confirm, as expected, that the asymptotic behavior of the oscillating scalar field is characterized by $\langle\gamma_\phi\rangle = 2n/(2+n)$. In (a) we show the time evolution of Ω_ϕ with a ϕ^2 potential ($n = 2$). Since the oscillating scalar field behaves like dust for $n = 2$, it gradually dominates the radiation and Ω_ϕ oscillates, asymptoting to unity. Similar behavior is seen in Ht in (b), where it approaches to $1/3$ corresponding to the cosmic expansion law of a dust dominated universe in the ρ^2 dominating era. In (c) we show the time evolution of Ω_ϕ for the ϕ^4 potential, ($n = 4$). Since the oscillating field behaves like radiation here, after a few oscillations, it approaches a constant value. In (d), for $n = 4$, $Ht \rightarrow \frac{1}{4}$, corresponding to the cosmic expansion law of a dust dominated universe in the ρ^2 dominated era.

has a distinct advantage in the preheating phase [44] if this scalar field couples to other matter fields. Since the cosmic expansion at late times becomes slower and the amplitude of ϕ decreases more slowly than in the conventional case, there may well be sufficient particle production generated as the field decays.

VII. SUMMARY

In this paper, we have studied the novel dynamics of a scalar field plus barotropic fluid, in a brane-world scenario dominated by the ρ^2 term in the effective four-dimensional Friedmann constraint. As a concrete example, we have adopted the Randall-Sundrum II model and

$\lambda = -\frac{\sqrt{2}m_5^{3/2}}{\sqrt{H}}\frac{V'}{V}$	$\Omega_\phi = \frac{\rho_\phi}{\rho}$	γ_ϕ	$V(\phi)$
$c = \text{cte} \left(> \sqrt{\frac{3\gamma}{2}} \right)$	$\frac{3\gamma}{2c^2}$	γ	$\mu^6\phi^{-2}$
$c = \text{cte} \left(< \sqrt{\frac{3\gamma}{2}} \right)$	1	$\frac{2c^2}{3}$	$\mu^6\phi^{-2}$
∞ (no oscil.)	0	γ	$\mu^{n+4}\phi^{-n}, n > 2$
∞ (oscil.)	0	$\frac{2n}{2+n} (> \gamma)$	$\frac{\mu^{4-n}}{n}\phi^n, n > 0, \text{ even}$
	cte	$\frac{2n}{2+n} (= \gamma)$	
	1	$\frac{2n}{2+n} (< \gamma)$	
0	1	0	$\mu^{n+4}\phi^{-n}, 2 > n > 0$

TABLE I: The asymptotic behavior of the contribution of the scalar field to the total energy density (Ω_ϕ) and the effective equation of state for the scalar field (γ_ϕ) for different limiting cases of $\lambda(\phi)$ in a ρ^2 dominated cosmology. In the last column we give an example of the potential $V(\phi)$ which satisfies this limit. Note how the duality transformation implies solutions match those in [1] for the variables λ , X and Y , although the definitions of those variables differ. In particular, the corresponding potentials in the last column differ for the two cases.

assumed that the scalar field is confined to our four dimensional spacetime.

Our approach has allowed us to deal with general classes of potentials, and complements an earlier investigation of a similar system but for particular potentials in [41]. Perhaps the most important result we have obtained, can be seen in the defining equations (10)-(19) in which we introduce a new set of variables to analyse the evolution equations in a model independent manner. The crucial point is that the equations for X and Y , given by (14) and (15) are identical to those derived in the conventional four dimensional cosmology where the Friedmann equation is driven by the energy density ρ of the scalar field and barotropic fluid, [1, 11, 14] as opposed to the ρ^2 driving term in our case. Although the precise definitions of X and Y differ, the fact that they obey the same evolution equation allows us to immediately write down and understand the form of the scaling solutions. There is a simple map which allows us to relate the cosmological solutions in each regime – a duality between the parameters X , Y in the two cases.

We have summarized our results in Table I, in an analogous manner to that presented in [1] for the equivalent conventional ρ dominated cosmology. In particular we showed that all the model dependence is given by $\lambda = -\frac{\sqrt{2}m_5^{3/2}}{\sqrt{H}}\frac{V'}{V}$ and the adiabatic index of the barotropic fluid γ . Scalar potentials that do not require introducing nonzero minima are classified into one of three different limiting cases by the asymptotic behavior of λ : λ goes to a finite constant, zero, or infinity. In the first case, Ω_ϕ approaches a finite constant (different from one or zero) depending on the value of λ . It is worth noting that in the conventional cosmology this happens in the model with an exponential potential, $V(\phi) = \mu^4 \exp(-\lambda_{\text{CLW}\kappa_4}\phi)$, while in the brane-world cosmology, it happens with an

inverse square potential model, $V(\phi) = \mu^6 \phi^{-2}$ – a reflection of the duality relating X , Y and X_{CLW} , Y_{CLW} .

In the second case $\lambda \rightarrow 0$, we obtained $X \rightarrow 0$, $Y \rightarrow 1$ corresponding to inflation with an almost constant Hubble parameter. As a concrete example, we considered the model with $V(\phi) = \mu^{4+n} \phi^{-n}$. For this potential, in the conventional cosmology, the inflationary solution is an attractor for any $n > 0$, but in the presence of the quadratic density term, it occurs only for $2 > n > 0$.

In the final case $\lambda \rightarrow \infty$. Following the procedure in [1] we investigated two different possibilities depending on whether λ oscillated or not. If it didn't, X , Y , $\Omega_\phi \rightarrow 0$ asymptotically. We showed the model with an exponential potential belongs to this class in the presence of the quadratic energy density term, as do models with inverse power law potentials with $n > 2$. As pointed out in [42], this provides a new feature for the quintessence scenario in that it allows us to explain how the energy densities in the radiation and scalar field could be different as we enter the usual ρ dominated Friedmann cosmology era. On the other hand, if λ does oscillate, the adiabatic index of the scalar field is given as $\langle \gamma_\phi \rangle = 2n/(2+n)$, where n is the power of the leading term in the scalar potential. The asymptotic behavior of the universe is decided by whether $\langle \gamma_\phi \rangle$ is larger than γ or not. For $\langle \gamma_\phi \rangle > \gamma$, $\Omega_\phi \rightarrow 0$, while for $\langle \gamma_\phi \rangle < \gamma$, $\Omega_\phi \rightarrow 1$ and for $\langle \gamma_\phi \rangle \gamma$, $\Omega_\phi \rightarrow cte$.

Although, we have concentrated on investigating the cosmology in the ρ^2 dominated regime, any realistic cosmology also has to allow for the fact that the universe is dominated today by the standard ρ term in the Fried-

mann equation. The next thing to do is to match the two regimes together. This has been investigated by a number of authors [34, 35, 36, 37, 38, 39, 40, 41] mainly using numerical simulations involving particular potentials. Such quintessential inflation models are being tightly constrained by recent CMBR data as they tend to generate too large a tensor contribution to the CMBR power spectra [35, 40]. What we have developed in this paper is an alternative, possibly powerful approach which allows us to make use of the duality that exists between the two regimes (ρ^2 and ρ dominated). We are currently investigating using this duality to determine consistent cosmologies involving an evolution from one regime into the other in a model independent manner, but based on the idea of using the definition of λ and γ as the key ingredients in the analysis.

Acknowledgments

S. M. would like to thank Kei-ichi Maeda for the continuous encouragement. He is grateful to the University of Sussex for their hospitality during a period when this work was initiated. S. J. L. would like to acknowledge the support of the Overseas Research Students Awards for financial support. This work was partially supported by The 21st century COE Program (Holistic Research and Education Centre for Physics self-organization systems at Waseda University.)

-
- [1] A. de la Macorra and G. Piccinelli, Phys. Rev. D **61**, 123503 (2000) [arXiv:hep-ph/9909459].
 - [2] A. Linde, *Particle Physics and Inflationary Cosmology*, (Harwood academic publishers, 1980); E. W. Kolb, and M. S. Turner, *The Early Universe*, (Perseus Publishing, 1990); A. R. Liddle, and D. H. Lyth, *Cosmological Inflation and Large-Scale Structure*, (Cambridge University Press, Cambridge, 2000).
 - [3] A. Vilenkin, Phys. Rept. **121**, 263 (1985); T. W. B. Kibble, J. Phys. **A9**, 1387 (1976); J. Preskill, Ann. Rev. Nucl. Part. Sci. **34**, 461 (1984).
 - [4] M. B. Hindmarsh and T. W. Kibble, "Cosmic strings," Rept. Prog. Phys. **58**, 477 (1995) [arXiv:hep-ph/9411342];
A. Vilenkin and E.P.S. Shellard, *Cosmic strings and other topological defects*, Cambridge Univ. Press (Cambridge 1994).
 - [5] M. B. Green, J. H. Schwarz and E. Witten, "Superstring Theory. Vol. 1 and 2", CUP. (1987)
 - [6] T. Barreiro, B. de Carlos and E. J. Copeland, Phys. Rev. D **58**, 083513 (1998) [arXiv:hep-th/9805005]; G. Huey, P. J. Steinhardt, B. A. Ovrut and D. Waldram, Phys. Lett. B **476**, 379 (2000) [arXiv:hep-th/0001112]; T. Barreiro, B. de Carlos and N. J. Nunes, Phys. Lett. B **497**, 136 (2001) [arXiv:hep-ph/0010102].
 - [7] B. Ratra and P. J. E. Peebles, Phys. Rev. D **37**, 3406 (1988).
 - [8] R. R. Caldwell, R. Dave and P. J. Steinhardt, Phys. Rev. Lett. **80** (1998) 1582 [arXiv:astro-ph/9708069]; I. Zlatev, L. M. Wang and P. J. Steinhardt, Phys. Rev. Lett. **82**, 896 (1999) [arXiv:astro-ph/9807002]; P. J. Steinhardt, L. M. Wang and I. Zlatev, Phys. Rev. D **59**, 123504 (1999) [arXiv:astro-ph/9812313].
 - [9] C. Wetterich, Nucl. Phys. **B302**, 668 (1988)
 - [10] P. G. Ferreira and M. Joyce, Phys. Rev. D **58**, 023503 (1998), **astro-ph/9711102**;
 - [11] E. J. Copeland, A. R. Liddle and D. Wands, Phys. Rev. D **57**, 4686 (1998) [arXiv:gr-qc/9711068].
 - [12] A. R. Liddle and R. J. Scherrer, Phys. Rev. D **59**, 023509 (1999) [arXiv:astro-ph/9809272].
 - [13] V. Sahni and A. Starobinsky, Int. J. Mod. Phys. D **9**, 373 (2000), **astro-ph/9904398**.
 - [14] S. C. C. Ng, N. J. Nunes and F. Rosati, Phys. Rev. D **64**, 083510 (2001) [arXiv:astro-ph/0107321].
 - [15] P. S. Corasaniti and E. J. Copeland, Phys. Rev. D **67**, 063521 (2003) [arXiv:astro-ph/0205544].
 - [16] V. A. Rubakov and M. E. Shaposhnikov, Towards A Solution To The Cosmological Constant Phys. Lett. B **125**, 139 (1983); K. Akama, Lect. Notes Phys. **176** (1982) 267 [arXiv:hep-th/0001113].
 - [17] P. Horava and E. Witten, Nucl. Phys. B **460**, 506 (1996)

- [arXiv:hep-th/9510209]; P. Horava and E. Witten, Nucl. Phys. B **475**, 94 (1996) [arXiv:hep-th/9603142].
- [18] A. Lukas, B. A. Ovrut and D. Waldram, Nucl. Phys. B **532**, 43 (1998) [arXiv:hep-th/9710208].
- [19] A. Lukas, B. A. Ovrut, K. S. Stelle and D. Waldram, Phys. Rev. D **59**, 086001 (1999) [arXiv:hep-th/9803235].
- [20] A. Lukas, B. A. Ovrut, K. S. Stelle and D. Waldram, Nucl. Phys. B **552**, 246 (1999) [arXiv:hep-th/9806051].
- [21] N. Arkani-Hamed, S. Dimopoulos and G. R. Dvali, Phys. Lett. B **429**, 263 (1998) [arXiv:hep-ph/9803315] I. Antoniadis, N. Arkani-Hamed, S. Dimopoulos and G. R. Dvali, Phys. Lett. B **436**, 257 (1998) [arXiv:hep-ph/9804398].
- [22] L. Randall and R. Sundrum, Phys. Rev. Lett. **83**, 3370 (1999) [arXiv:hep-ph/9905221]; L. Randall and R. Sundrum, Phys. Rev. Lett. **83**, 4690 (1999) [arXiv:hep-th/9906064].
- [23] A. Lukas, B. A. Ovrut and D. Waldram, arXiv:hep-th/9802041.
- [24] A. Lukas, B. A. Ovrut and D. Waldram, Phys. Rev. D **61**, 023506 (2000) [arXiv:hep-th/9902071].
- [25] E. J. Copeland, J. Gray and A. Lukas, Phys. Rev. D **64**, 126003 (2001) [arXiv:hep-th/0106285].
- [26] R. Maartens, arXiv:gr-qc/0101059; R. Maartens, Prog. Theor. Phys. Suppl. **148**, 213 (2003) [arXiv:gr-qc/0304089]; R. Maartens, arXiv:gr-qc/0312059.
- [27] D. Langlois, arXiv:gr-qc/0207047; D. Langlois, Prog. Theor. Phys. Suppl. **148**, 181 (2003) [arXiv:hep-th/0209261].
- [28] P. Brax and C. van de Bruck, Class. Quant. Grav. **20**, R201 (2003) [arXiv:hep-th/0303095]; P. Brax, C. van de Bruck and A. C. Davis, arXiv:hep-th/0404011.
- [29] D. Wands, Class. Quant. Grav. **19**, 3403 (2002) [arXiv:hep-th/0203107]; E. Papantonopoulos, Lect. Notes Phys. **592**, 458 (2002) [arXiv:hep-th/0202044]; G. Gabadadze, arXiv:hep-ph/0308112.
- [30] T. Shiromizu, K. i. Maeda and M. Sasaki, Phys. Rev. D **62**, 024012 (2000) [arXiv:gr-qc/9910076].
- [31] P. Binetruy, C. Deffayet and D. Langlois, Nucl. Phys. B **565**, 269 (2000) [arXiv:hep-th/9905012] P. Binetruy, C. Deffayet, U. Ellwanger and D. Langlois, Phys. Lett. B **477**, 285 (2000) [arXiv:hep-th/9910219].
- [32] S. Mukohyama, T. Shiromizu and K. i. Maeda, Phys. Rev. D **62**, 024028 (2000) [Erratum-ibid. D **63**, 029901 (2001)] [arXiv:hep-th/9912287].
- [33] K. Ichiki, M. Yahiro, T. Kajino, M. Orito and G. J. Mathews, Phys. Rev. D **66**, 043521 (2002) [arXiv:astro-ph/0203272]; J. D. Bratt, A. C. Gault, R. J. Scherrer and T. P. Walker, Phys. Lett. B **546**, 19 (2002) [arXiv:astro-ph/0208133].
- [34] E. J. Copeland, A. R. Liddle and J. E. Lidsey, Phys. Rev. D **64**, 023509 (2001) [arXiv:astro-ph/0006421]; G. Huey and J. E. Lidsey, Phys. Lett. B **514**, 217 (2001) [arXiv:astro-ph/0104006].
- [35] V. Sahni, M. Sami and T. Souradeep, Phys. Rev. D **65**, 023518 (2002) [arXiv:gr-qc/0105121]; M. Sami and V. Sahni, arXiv:hep-th/0402086.
- [36] A. S. Majumdar, Phys. Rev. D **64**, 083503 (2001) [arXiv:astro-ph/0105518].
- [37] N. J. Nunes and E. J. Copeland, Phys. Rev. D **66**, 043524 (2002) [arXiv:astro-ph/0204115].
- [38] J. E. Lidsey and N. J. Nunes, Phys. Rev. D **67**, 103510 (2003) [arXiv:astro-ph/0303168].
- [39] S. Tsujikawa and A. R. Liddle, JCAP **0403**, 001 (2004) [arXiv:astro-ph/0312162].
- [40] M. Sami and N. Dadhich, arXiv:hep-th/0405016.
- [41] S. Mizuno, K. i. Maeda and K. Yamamoto, Phys. Rev. D **67**, 023516 (2003) [arXiv:hep-ph/0205292].
- [42] K. i. Maeda, Phys. Rev. D **64**, 123525 (2001) [arXiv:astro-ph/0012313]; S. Mizuno and K. i. Maeda, Phys. Rev. D **64**, 123521 (2001) [arXiv:hep-ph/0108012].
- [43] K. i. Maeda and D. Wands, Phys. Rev. D **62**, 124009 (2000) [arXiv:hep-th/0008188].
- [44] S. Tsujikawa, K. i. Maeda and S. Mizuno, Phys. Rev. D **63**, 123511 (2001) [arXiv:hep-ph/0012141].
- [45] F. Lucchin and S. Matarrese, Phys. Rev. D **32**, 1316 (1985).
- [46] P. J. E. Peebles and A. Vilenkin, Phys. Rev. D **59**, 063505 (1999) [arXiv:astro-ph/9810509].